What Is COVID-19?

Coronavirus disease 2019 (COVID-19) is a disease caused by a new coronavirus, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2).

Coronaviruses are a cause of the common cold, but SARS-CoV-2, which likely came from bats, causes more severe disease in many patients.¹

Symptoms and Diagnosis

Symptoms occur on average about 5 days after exposure to the virus. Almost all patients develop symptoms within 12 days. Therefore, a 14-day self-isolation is recommended for people who were likely exposed to the virus. The most common symptoms are cough, fever, and shortness of breath; most patients with COVID-19 have at least 1 of these. Other common symptoms include muscle aches, fatigue, nausea, vomiting, and diarrhea. People may also have loss or change in sense of taste and smell.²

COVID-19 is typically diagnosed by a swab of the nose that measures the coronavirus. There are also blood tests that can measure recent exposure to the virus, but these do not show positive results until 1 to 3 weeks after infection starts.

COVID-19 Disease Progression and Treatment

The virus typically enters through the eyes, mouth, or nose then travels down the throat, where it may cause a cough. In some patients, the virus enters the lungs and may cause pneumonia. Pneumonia leads to fluid filling the air sacs in the lungs, which makes breathing difficult. Most patients with pneumonia must be hospitalized and treated with oxygen. Some patients become very ill and need life support such as mechanical ventilation. About 1 in 20 patients with COVID-19 dies. However, death rates vary substantially by age, ranging from 1 in 900 patients aged 18 to 29 years to 1 in 34 aged 50 to 64 years and 1 in 3 aged 85 years or older.

There is currently no vaccine to reduce risk of COVID-19. Several potential vaccines are in development, which may help prevent COVID-19 in the future. For most patients, treatment is supportive. Antibiotics do not work for this viral disease. Studies are testing many antiviral medications, as well as medications to modify the body's response to the virus. For certain hospitalized patients, antiviral drugs and steroids may help.


Authors: W. Joost Wiersinga, MD, PhD, MBA; Hallie C. Prescott, MD, MSc Published Online: July 10, 2020. doi:10.1001/jama.2020.12984

Author Affiliations: Division of Infectious Diseases, Department of Medicine, Amsterdam UMC, Location UMC, University of Amsterdam, Amsterdam, the Netherlands (Wiersinga); Division of Pulmonary and Critical Care Medicine, University of Michigan, Ann Arbor (Prescott).

Conflict of Interest Disclosures: Dr Wiersinga reported receiving grants from the Netherlands Organisation of Scientific Research. Dr Prescott reported receiving grants from the National Institutes of Health, Agency for Healthcare Research and Quality, and Department of Veterans Affairs and serving on the Surviving Sepsis Campaign.

1. Wiersinga WJ, Rhodes A, Cheng AC, et al. Pathophysiology, transmission, diagnosis, and treatment of coronavirus disease 2019 (COVID-19): a review. JAMA. Published online July 10, 2020. doi: 10.1001/jama.2020.12839 2. Huynh PP, Ishii LE, Ishii M. What is anosmia? JAMA. Published online June 18, 2020. doi:10.1001/jama. 2020.10966





Patients with COVID-19 are likely to have fatigue and decreased energy for 6 to 8 weeks. It may be difficult to resume the same activity and exercise as before COVID-19, but it is important to remain active and increase exercise gradually. For most patients, lung function returns to normal after pneumonia.

Scientists do not know if people can have COVID-19 more than once, but repeat infection may be possible, particularly for people with mild symptoms during their initial infection.

FOR MORE INFORMATION

JAMA COVID-19 Information Page jamanetwork.com/journals/jama/pages/coronavirus-alert

The JAMA Patient Page is a public service of JAMA. The information and recommendations appearing on this page are appropriate in most instances, but they are not a substitute for medical diagnosis. For specific information concerning your personal medical condition, JAMA suggests that you consult your physician. This page may be photocopied noncommercially by physicians and other health care professionals to share with patients. To purchase bulk reprints, email reprints@ iamanetwork.com.